

## FlyCut™ XhoI

Cat.No. JX201

Storage: at -20°C for two years

Concentration: 20,000 units/ml

### Description

*FlyCut*™ XhoI is expressed and purified from *E.coli* that carries the recombinant XhoI gene. The molecular weight is 27.9 kDa, with the recognition site at C<sup>^</sup>TCGAG. The reaction is conducted for 5-10 minutes at 37°C, and heat-inactivated at 65°C for 20 minutes. This enzyme is not sensitive to dam or dcm methylation, but sensitive to mammalian CpG methylation.

### Enzyme Properties

Fast digestion in 5-10 minutes with high fidelity

### Application

Genomic DNA, plasmid DNA, PCR product

### Kit Contents

Component	JX201-01	JX201-02
<i>FlyCut</i> ™ XhoI	2,500 units	5,000 units
10× <i>FlyCut</i> ™ Buffer	1 ml	1 ml
10×DNA Loading Buffer	1 ml	1 ml

### Unit Definition

One unit is defined as the amount of enzyme required to digest 1 µg of λ DNA in 1 hour at 37°C in a total reaction volume of 50 µl.

### Quality Control

**Ligation and re-cutting:** After 10-fold overdigestion with *FlyCut*™ XhoI, more than 95% of the DNA fragments can be ligated with T4 DNA ligase at 25°C. Of these ligated fragments, more than 95% can be recut.

**16-Hour incubation:** A 50 µl reaction containing 1 µg of DNA and 10 units of enzyme incubated for 16 hours results in the same pattern of DNA bands as a reaction incubated for 1 hour with 1 unit of enzyme.

**Blue/White screening (Terminal integrity):** A DNA vector is digested at a unique site within lacZα gene with a 10-fold excess of enzyme, and then ligated, transformed and plated on X-gal/IPTG plate. Successful expression of the β-galactosidase indicates that lacZα gene remains integrity after cloning. A blue colony represents an intact gene, and a white colony represents an interrupted gene. To be Blue/White certified, enzymes must produce fewer than 3% white colonies.

**Exonuclease activity:** After incubation for 4 hours at 37°C, a 50 µl reaction containing 100 units of enzyme and 1 µg <sup>3</sup>H DNA releases less than 0.1% radioactive substance.

**Endonuclease activity:** After incubation for 4 hours at 37°C, a 50 µl reaction containing 15 units of enzyme with 1 µg pBR322 RFI DNA results in less than 10% conversion from RFI to RFII.

### Storage Buffer

20 mM Tris-HCl pH7.4, 250 mM NaCl, 0.1 mM EDTA, 1.5 mM DTT, 400 µg/ml BSA, 50% Glycerol

### 10×*FlyCut*™ Buffer

500 mM Tris-Ac pH7.9, 1 M KAc, 120 mM MgAc<sub>2</sub>, 1 mg/ml BSA

### Reaction Components

Component	20 µl Volume	50 µl Volume
DNA	≤1 µg	≤2 µg
10× <i>FlyCut</i> ™ Buffer	2 µl	5 µl
<i>FlyCut</i> ™ XhoI	0.5 µl	1 µl
ddH <sub>2</sub> O	to 20 µl	to 50 µl



**Trans, The Highest Quality  
Products For Life Sciences**

Prior to use, please completely mix the *FlyCut*<sup>™</sup> Buffer. Increase the volume of enzyme, in case of digestion of >2 µg DNA or incomplete digestion, but the total volume of enzyme should be less than 1/10 of the reaction system. Incubation for 5-10 minutes at 37°C. Enzyme is inactivated by adding 10×DNA Loading Buffer to a final concentration at 1×, or by heating at 65°C for 20 minutes.

**Notes**

- Thaw the 10×*FlyCut*<sup>™</sup> Buffer completely and mix well before use.
- Low ionic strength, high enzyme concentration, glycerol concentration > 5%, or pH > 8.0 may result in star activity.

**FOR RESEARCH USE ONLY**